

**ANTI-AGEING ACTIVITY AND IDENTIFICATION OF BIO ACTIVE COMPOUND
FROM THE SEED OF *Linum usitatissimum* BY GC-MS TECHNIQUES**

S.REXCIDA JANTHARK MARY*, S.JOSEPHINOL¹, B. RAMYA² AND SEBATINA LOUIS³

Department of Biochemistry, Holy Cross College (Autonomous), Tiruchirappalli, India

*Corresponding Author: S. Rexcida Janthark Mary: E Mail: rexidadepthi@gmail.com

Received 21st March 2021; Revised 25th April 2021; Accepted 12th June 2021; Available online 1st Feb. 2022

<https://doi.org/10.31032/IJBPAS/2022/11.2.5894>

ABSTRACT

Introduction: Aging occurs in all living organisms and it is an inevitable process. Aging is caused due to both intrinsic and extrinsic factors. Intrinsic aging is an inevitable physiological process. Extrinsic aging occurs in addition to intrinsic aging as a result of sun and environmental damage. Activation of hyaluronidase, collagenase and elastase, leads to skin aging. By using the medicinal crop *Linum usitatissimum* were investigated to assess their skin aging properties.

Methods: The activity of anti- elastase, anti- collagenase and anti-hyaluronidase in ethanol extract from the seed of *Linum usitatissimum* was determined by using spectrophotometric methods. The bio-active compounds of *Lainum usitatissimum* was analyzed using GC-MS method.

Result: The anti-elastase assay in ethanol extract of *Linum usitatissimum* shows 70% inhibition, for anti- collagenase assay shows 73% inhibition and for anti-hyaluronidase assay shows 61% inhibition. The ethanol extract of *Linum usitatissimum* contained the highest anti-collagenase activity.

Conclusion: The enzymes which are responsible for aging is inhibited by ethanol extract of *Linum usitatissimum*, suggests that it can help to restore skin elasticity and slows down the process of aging.

Keywords: Anti-ageing Activity, *Linum usitatissimum*, Spectrophotometric method, GC-MS Techniques

INTRODUCTION

Aging occurs in all living organisms and it is an inevitable process. The skin protects us from microbes and the elements, helps to regulate body temperature, and permits the sensation of touch, heat and cold. Chronological aging and photoaging are the two types of skin aging [1]. The extrinsic factor causes photoaging which includes symptoms like leathery appearance, dark or light pigmentation and deep furrows [2, 3]. The epidermis, dermis and subcutaneous tissue are the three layers in skin [4]. The outermost part of the skin is the extracellular matrix (ECM) which is made up of fibroblasts and proteins that include collagens and elastin [5]. The ECM plays an important role in the maintenance of physiological functions of the body because it is essential for growth and elasticity of the skin [6]. Skin aging is directly linked with degradation of ECM and it is correlated with an increase in activity of some enzymes, which is involved in skin aging including hyaluronidase, collagenase and elastase [3, 7, 8].

The main component of connective tissue, hair and nails is collagen and is the building blocks of skin [1]. The elasticity, strength and flexibility of the skin is maintained by collagen. The moisture of the skin is retained by hyaluronic acid and it is also responsible for its structure and elasticity of the skin. It is involved in rapid tissue

proliferation, regeneration and repair, it also facilitates the exchange of nutrients and base products [9, 10]. Hyaluronic acid is required for the organization and structural maintenance of the ECM. When collagen, elastin and hyaluronic acid level decreases it leads to loss of strength and flexibility in the skin, which further results in visible wrinkles.

Linum usitatissimum Linn, commonly known as Alsi belongs to the family Linaceae and it is cultivated in cooler regions throughout the world [11]. It has anticancer [12, 13], antidiabetic [14], antimicrobial [15], analgesic, antipyretic and anti-inflammatory activity [16], and it helps to reduce cardiovascular diseases [17]. Its oil is known as linseed oil. It is used for various medicinal purposes. Textiles made from flax are known as linen and are used for bed sheets, underclothes and table linen [18]. This flax seed protein helps to improve immune function, lowers cholesterol, prevents tumor, has antifungal activity and it has been proved by animal studies [19-21].

MATERIALS AND METHODS

Collection of plant materials and preparation of ethanolic extracts:

The *Linum usitatissimum* stem seed is collected from the nearby villages in trichy and confirmed its presence with

the help of guide. In a uniform powdered manner of 40 mesh size the *Linum usitatissimum* seed are grinded, before grinding the *Linum usitatissimum* seed is washed in water and dried it for a week and place it in the room temperature. The 20g of dried seed powder of *usitatissimum* is dissolved in a prepared ethanolic extract (40ml) and keep it in a hot percolation for 1 hour of duration. After cooling the extract, it is filtered by using Whatmann filter paper No.42 (125mm). By using this method the undissolved products are removed which contain cellular material and some other materials. The solution which is filtered is used for the further analysis of qualitative and quantitative method.

Chemicals and reagents:

FALGPA [N-[3-(2-furyl)acryloyl]-leu-Gly-Pro-Ala, TES[tris(hydroxymethyl)-methyl-2-aminoethane sulphonate (TES), The type I collagenase from clostridium, histolytium, ethanol, isopropanol, ninhydrine solution, citrate buffer, 4-dimethylamino benzaldehyde (DMAB), The HEPES at pH 7.5, 4-(2-hydroxyethyl)-1-piperazine ethanesulfonic acid, Potassium metaborate (KBO₂), Hyaluronic acid, Hyaluronidase acid, calcium chloride dihydrate, human leukocyte elastase, N-methoxysuccinyl-Ala-Ala-Pro-Chloro.

The anti-elastase activity determination:

For the anti-elastase activity determination, the preparation for bulk needs various concentration of *Linum usitatissimum* seed extract (20,40,60,80,100 µg/ml) was taken in the test tube and labelled it; the 100µg/ml of extract is added with 900 µg/ml of ethanol and 200 µg/ml extract is added with 800 µg/ml of ethanol and 300 µg/ml extract added with 700 µg/ml of ethanol, 400 µg/ml of extract is added with 600 µg/ml of ethanol, and 500 µg/ml of extract is added with 500 µg/ml of ethanol. For the control solution ethanol (1000µg/ml) is taken. The elastase of 500µg/ml is taken and added with 500µg/ml hepes buffer and then add 1000µg/ml of various concentration of plant sample extract and keep it in room temperature for 20 minutes then add 500µg/ml N-methoxysuccinyl-ala-ala-pro-chloro in test tubes which are incubated for further 20 minutes then check the absorbance at 540nm. The percentage inhibition was calculated as follows:

$$\text{Inhibition (\%)} = \frac{(\text{A control} - \text{A sample})}{\text{A control}} \times 100$$

Anti-hyaluronidase activity determination:

The method of finding out hyaluronidase activity is based on flurometric Morgan - Elson assay

Method. 500µg/ml calcium chloride which was taken and added it with 1000µg/ml of various

Concentration from the bulk solution and then add 500µg/ml hyaluronidase acid keep it for 20 minutes in a room temperature. Then add 250µg/ml of hyaluronic acid they are all kept in it for 10 minutes. The 250µg/ml of KBO₂ is added in a test tube. They are heated and cooled for 5 minutes then 500µg/ml DNAB is added in test tube keep it for 10 minutes incubation in room temperature check for the absorbance at 540 nm.

Anti- collagenase activity determination:

The method for finding out anti-collagenase activity is based on the Moore stein method which was modified by Mandal et.al. The 500µg/ml of collagenase is taken in a test tube with that 500µg/ml of TRIS buffer at the various concentration of the extract 1000 µg/ml is taken. They are all kept for 10 minutes incubation in a room temperature and then add 500µg/ml FALGPA enzyme which has been keep it in room temperature for 10 minutes then 500 µg/ml of citrate buffer solution is added and incubate it for 5 minutes in room temperature. 500µg/ml ninhydrin solution is added heat and cool it for 5 minutes and then 500µg/ml isopropanol which is added and incubate it for 5 minutes and check for the absorbance at 540nm.

GC-MS Analysis of Flax Seed:

GC/MS analysis was performed at Heber Analytical Instrumentation Facility, Thiruchirapalli, for MS identification of the

GC components. The column used was DB-5 (J & W Scientific, Folosm, CA) cross-linked fused silica capillary column (30 m long, 0.25 mm internal diameter) coated with polydimethylsiloxane (0.5µm film thickness). The oven temperature was programmed from 50oC for 2 min., at isothermal, then heating by 7oC/ min. to 250oC and isothermally for 10 min., at 250oC. Injector temperature was 250oC and the volume injected was 0.5 µl. Transition line and ion source temperature were 250oC and 200oC respectively. The mass spectrometer had a delay of 2 min. to avoid the solvent peak and then scanned from m/z 50 to m/z 500. Ionization energy was set at 70 eV.

RESULTS AND DISCUSSIONS**Anti - elastase activity determination:**

In the place of connective tissues elastin is found. The elastin provides the elasticity to our skin and lungs. It is a kind of protein and it is monitorized by the enzyme called elastase. If the elastin content in our body is decreased or by exposure to UV radiation causes skin aging. Our crop *Linum usitatissimum* has provided a good firm to our skin and improve the elasticity of our skin. The ethanolic extract of seed of *Linum usitatissimum* at different varying concentrations (100µg/ml, 200µg/ml, 300µg/ml, 400µg/ml, 500µg/ml) shows the various % inhibition concentration and the maximum inhibition concentration is

showed at 100µg/ml =53.19% which is compared with the standard Elafin which show the % concentration at 100µg/ml=48.23% (Table 1, Figure 1).

Anti collagenase activity determination:

The skin is the most important layer in our body and it consist of some important components like collagen with it. The collagen is declined by the presence of enzyme collagenase. If the collagenase is hindered means the aging process slowly occurs and the formation of pre-collagen fibres is delayed. The ethanolic extract of seed of *Linum usitatissimum* with different varying concentration (109µg/ml, 200µg/ml, 300µg/ml, 4000µg/ml, 500µg/ml) shows the different % inhibition concentration and the maximum % inhibition concentration in the anti-collagenase activity shows at 100µg/ml=56.25% and it is compared to the standard EDTA which show the anti-collagenase Activity at 100µg/ml=73.44% (Table 2, Figure 2).

Anti-hyaluronidase activity:

The hyaluronidase activity is more important for our skin because it reduces the fine lines in our skin and it consists of some healing property with it. The ethanolic extract of the seed of

usitatissimum in the different concentrations (100µg/ml, 200µg/ml, 300µg/ml, 400µg/ml, 500µg/ml) show the different % inhibition concentration and the maximum % inhibition concentration is shown at the range of 100µg/ml=39.81% which is compared to the standard Sodium aurothiomalate. To find out the anti-hyaluronidase activity in the standard the maximum % inhibition concentration is shown at 100µg/ml= 74.07% (Table 3 and Figure 3).

Gas chromatography- mass spectrometry of flax seed:

Bicyclo[4.1.0]Hept-3-Ene-2-Thiol,3,7,7-Trimethyl-,[1S-(1.Alpha.,2.Alpha.,6.Alpha.)]-; Heptadecanoic Acid, Methyl Ester; Hexadecanoic Acid; Methyl Octadeca-9,12-Dienoate; (Z,Z) - 6,9-CIS-3,4-Epoxy-Nonadecadiene; 9,12-Octadecadienoic Acid, Methyl Ester, (E,E)-; 7-Tetradecenal, (Z)-; Bicyclo[3.3.1]nonan-1-ol; 9-Octadecenoic Acid (Z)-; 1-Naphthalenemethanol, 1,4,4a,5,6,7,8,8a-Octahydro-2,5,5,8a-Tetramethyl-; 1-Undecene-5,9-Diyne; 1,3,5-Triphenyl-1,5-Pentanedione; Silane, methyl-di-isopropoxyethoxy-; (+)-2,3-o-Benzylidene-D-threitol; Chromium, (.eta.-5-cyclopentadienyl)-(.eta.-6-toluene)

(Table 4, Figure 4).

Table 1: Determination of anti-elastase activity of seed of *Linum usitatissimum* with the standard Elafin

CONCENTRATION	ANTI - ELASTASE % inhibition concentration	
	Seed of <i>Linum usitatissimum</i>	Elafin
100µg/ml	53.19	67.37
200µg/ml	58.16	73.04
300µg/ml	61.70	78.01

400µg/ml	65.96	82.26
500µg/ml	70.92	85.81




Figure 1: Comparative Graph of Seed *Linum usitatissimum* with the Standard Elafin

Table 2: Anti-collagenase activity of seed of *Linum usitatissimum* with the standard EDTA

Concentration	Anti-collagenase % inhibition	
	Seed of <i>usitatissimum</i>	EDTA
100µg/ml	56.25	73.43
200µg/ml	62.5	77.34
300µg/ml	67.18	79.68
400µg/ml	69.53	82.03
500µg/ml	73.43	84.37




Figure 2: Comparative graph of *Linum usitatissimum* seed with the standard EDTA

Table 3: Anti-hyaluronidase activity of seed of *Linus usitatissimum* compared with sodium aurothiomalate

Concentration	Anti-hyaluronidase activity % inhibition concentration	
	Seed of <i>Linum usitatissimum</i>	Sodium aurothiomalate
100µg/ml	39.81	74.07
200µ/ml	45.37	77.77
300µg/ml	50.92	80.55
400µg/ml	55.55	83.33
500µg/ml	61.11	86.11









Figure 3: Comparative graph on seed of *Linum usitatissimum* with the standard Sodium aurothiomalate

Figure 4: Gas Chromatography- Mass Spectrometry of Flax Seed

Table 4: Gas Chromatography- Mass Spectrometry of Flax Seed

COMPOUND NAME	RETENTION	AREA%	MOLECULAR FORMULAE	MOLECULAR STRUCTURE
Bicyclo [4.1.0] Hept-3-Ene-2-Thiol, 3,7,7-Trimethyl-, (1. Alpha., 2. Alpha., 6. Alpha.)-	7.401	1.33	C ₁₀ H ₁₆ S	
Heptadecanoic Acid, Methyl Ester	26.012	2.54	C ₁₈ H ₃₆ O ₂	
Hexadecanoic Acid	26.613	8.14	C ₁₆ H ₃₂ O ₂	
Methyl Octadeca-9,12-Dienoate	28.683	6.66	C ₁₉ H ₃₄ O ₂	
(Z,Z)-6,9-Cis-3,4-Epoxy-Nonadecadiene	28.779	11.02	C ₁₉ H ₃₄ O	
9,12-Octadecadienoic Acid, Methyl Ester, (E,E)-	29.301	6.25	C ₁₉ H ₃₄ O ₂	
7-Tetradecenal, (Z)-	29.392	43.71	C ₁₄ H ₂₆ O	

Bicyclo[3.3.1]Nonan-1-Ol	29.62	2.76	C ₉ H ₁₆ O	
9-Octadecenoic Acid (Z)-	29.759	4.73	C ₁₈ H ₃₄ O ₂	
1-Naphthalenemethanol, 1,4,4a,5,6,7,8,8a-Octahydro-2,5,5,8a-Tetramethyl-	38.565	3.1	C ₁₅ H ₂₆ O	
1-Undecene-5,9-Diyne	38.614	1.81	C ₁₁ H ₁₄	
1,3,5-Triphenyl-1,5-Pentanedione	38.65	2	C ₂₃ H ₂₀ O ₂	
Silane, methyl-diisopropoxyethoxy-	38.675	2.5	C ₉ H ₂₂ O ₃ Si	
(+)-2,3-o-Benzylidene-D-threitol	39.29	1.72	C ₁₁ H ₁₄ O ₄	
Chromium, (eta-5-cyclopentadienyl)-(eta-6-toluene)	39.35	1.75	C ₁₂ H ₁₃ Cr	

CONCLUSION

The present study concluded with the facts that the fibre crop *usitatissimum* of *Linum usitatissimum* has the presence of the antiaging property and it is resulted under different methods. From the qualitative test it is confirmed that the seed of *Linum usitatissimum* in ethanolic extract has shown the presence of Tannin, Steroids, Saponin, Flavonoids, Alkaloids, Terpenoids, Phlobatannin, Phenol, Glucoanthocyanin, Cardiac glycosides, Anthocyanin,

Antraquinones, Glycosides, Coumarine, Emodin, Xanthoprotein, Carbohydrates and Protein content. The seed of *Linum usitatissimum* used for the treatment of anti-aging which is due to the presence of various phytochemicals component present in it.

The anti-elastase activity in the ethanolic extract of seed of *Linum usitatissimum* shown the maximum % inhibition concentration at 100µg/ml = 53.19% which

is compared to the standard elafin at 100µg/ml=48.23%. The anti-collagenase activity in the ethanolic extract of seed of *Linum usitatissimum* shown the maximum % inhibition concentration at 100µg/ml = 56.25% which is compared to the standard EDTA at 100µg/ml=73.44%. The anti-hyaluronidase activity in the ethanolic extract of seed of *Linum usitatissimum* which show the maximum % inhibition concentration at 100µg/ml = 39.81% which is compared to the standard sodium aurothiomalate at 100µg/ml = 74.07% so, it is confirmed that the seed of *Linum usitatissimum* is helpful for the treatment anti-aging.

From analyzing the above results the natural fibrecrop seed of *Linum usitatissimum* has confirmed its presence in the treatment of anti-aging.

ACKNOWLEDGEMENT

S. Rexcida Janthark Mary acknowledges Dr. K. Manjula, Chief Executive Officer, Bio Techno Solutions Training and Research Institute, Ponmalaipatti, Trichy for her constant support for this research.

REFERENCE

- [1] Mukherjee PK, Maity N, Nema NK, Sarkar BK Bioactive compounds from natural resources against skin aging, *Phytomedicine* 2011, 19:64-73.
- [2] Fisher GJ, Karni S, Varani J, Bata-Csorgo Z, Wan J, Data S, Voorhees JJ: Mechanisms of photoaging and chronological skin aging. *Arch Dermatol* 2002. 138(11):1462-1470.
- [3] Maity N, Nema NK, Abedy S, Sarkar BK, Mukherjee PK Exploring *Tagetes erecta* Linn flower for the elastase, hyaluronidase and MMP-1 inhibitory activity. *J Ethnopharmacol* 2011, 137:1300-1305.
- [4] Rittie L, Fisher G UV-light-induced signal cascades and skin aging. *Ageing Res Rev* 2002. 1:705-720.
- [5] Fulop T, Khalil A, Larbi A: The role of elastin peptides in modulating the immune response in aging and age-related diseases, *PatholBiol* 2012 60:28-33.
- [6] Kurtz A, Oh S- Age related changes of the extracellular matrix and stem cell maintenance. *Prev Med* 2012. 54: 550-556.
- [7] Wary KK, Thakkar GD, Humtsoe J, Yang Y Analysis of VEGF-responsive genes involved in the activation of endothelial cells. *Mol. Concer* 2003 2: 25-36.
- [8] Losses JN, Munene CN, Bansode RR, Bawadi HA: Inhibition of matrix metalloproteinase-1 activity by the soybean Bowman-Birk inhibitor. *BiotechnolLett* 2004, 26: 901-905.

- [9] Manuskiatti W, Maibach H: Hyaluronic acid and skin: wound healing and aging. *Int J Dermatol* 1996, 35(8): 539-544.
- [10] Hsu M-F, Chiang B-H: Stimulating effects of *Bacillus subtilis* fermented *Radix astragali* on hyaluronic acid production in human skin cells. *J Ethnopharmacol* 2009, 125: 474-481.
- [11] Shweta Gokhale, Alakh. N Sahu. Pharmacological properties of flaxseed, *Linum usitatissimum* Linn., as a potential medicinal plant: An overview. *World Journal of Pharmaceutical Sciences* 2016
- [12] Kuijsten A, Hollman PCH, Boshuizen HC, Buijsman MNC, Veer PV, Kok FJ *et al.* Plasma enterolignan concentrations and colorectal cancer risk in a nested casecontrol study. *Am. J. Epidemiol.* 2008; 167(6): 734-742.
- [13] Yan L, Yee JA, Li D, Mcguire MH, Thompson LU. Dietary flaxseed supplementation and experimental metastasis of melanoma cells in mice. *Cancer Letter.* 1998; 124(2):181-186.
- [14] Ghule AE, Jadhav SS, Bodhankar SL. Effect of ethanolic extract of seeds of *Linum usitatissimum* Linn. In hyperglycaemia associated ROS production in PBMNCs and pancreatic tissue of alloxan induced diabetic rats. *Asian Pacific Journal of Tropical Disease*, 2012, 405-410.
- [15] Amin T, Thakur M. A comparative study on proximate composition, phytochemical screening, antioxidant and antimicrobial activities of *Linum usitatissimum* L. (flax seeds). *Int J Curr Microbial App Sci.* 2014; 3(4):465- 481.
- [16] Kaithwas G, Mukherjee A, Chaurasia AK, Majumdar DK. Anti-inflammatory, analgesic and antipyretic activities of *Linum usitatissimum* L. (flaxseed/linseed) fixed oil. *Indian J Exp Biol.* 2011 Dec; 49(12): 932-8. PMID: 22403867.
- [17] Cintra DEC, Costa AGV, Peluzio MCG, Matta SLP, Silva MTC, Costa NMB. Lipid profile of rats fed high-fat diets based on flaxseed, peanut, trout, or chicken skin. *Nutrition.* 2006; 22(2):197-205.
- [18] Gambus H, Mikulec A, Gambus F, Pisulewski P. Perspectives of linseed utilization in baking. *Pol J Food Nutr Sci.* 2004; 13(54): 21-27.

- [19] Rabetafika, H.N., Van Remoortel, V., Danthine, S., Paquot, M. and Blecker, C. (2011), Flaxseed proteins: food uses and health benefits. *International Journal of Food Science & Technology*, 46: 221-228.
- [20] Verma A, Houtem KRJ, Raja, Sen S, Sachan S, Mishra A, pharmacological evaluation of *Saracaindica* leaves for CNS depressant activity in mice *J Pharm sci, Res.* vol. 2010, 2(6), 338- 343.
- [21] Xu Y, Hall C 3rd, Wolf-Hall C. Antifungal activity stability of flaxseed protein extract using response surface methodology. *J Food Sci.* 2008 Jan; 73(1): M9-14. [10.1111/j.1750-3841.2007.00576.x](https://doi.org/10.1111/j.1750-3841.2007.00576.x). PMID: 18211360.