



PHYTOCHEMICAL EVALUATION OF ROOTS OF *PLUMBAGO ZEYLANICA* AND ASSESSMENT OF ITS ANTIOXIDANT POTENTIAL

USMAN FN^{1*} AND SINGHAL NK²

1: Research Scholar, Department of Pharmacy, Shyam University, Dausa-303511, Rajasthan, India

2: Professor, Department of Pharmacy, Shyam University, Dausa-303511, Rajasthan, India

*Corresponding Author: Ms. Fathima Nausheen Usman: E Mail: bandameedi.ramu@gmail.com

Received 24th Sept. 2023; Revised 25th Oct. 2023; Accepted 12th Jan. 2024; Available online 1st Nov. 2024

<https://doi.org/10.31032/IJBPAS/2024/13.11.8385>

ABSTRACT

Plumbago zeylanica (Plumbaginaceae) commonly cultivated in India for its therapeutic value. The goal of current study was to evaluate the phytochemical constituents and to explore the in vitro antioxidant properties of ethanolic extracts of *P. zeylanica*. Preliminary phytochemical evaluations revealed the presence of steroids, tannins, glycosides, phenols, flavonoids and saponins in the Leaves methanolic extracts. Determinations of their in vitro antioxidant activity were carried out by using 2,2-diphenyl-1-picrylhydrazyl in an assay based on the method of Oyaizu. Hydroxyl radical scavenging activity was also measured in the same extracts. Results from this research project indicated that *P. zeylanica* Leaves are indeed rich in phytochemicals and had substantial antioxidant activities, implying that *P. zeylanica* Leaves can be used as a potential source of natural antioxidants.

Keywords: *Plumbago zeylanica*, Medicinal plants, Phytochemical screening, Antioxidant, DPPH assay, Reactive oxygen species

1. INTRODUCTION

Medicinal plants have been utilized for thousands of years as major sources of cures for human diseases [1]. Isolation and characterization of pharmacologically active compounds from medicinal plants have

gained vital importance in developing countries. More recently, drug discovery techniques have been applied to the standardization of herbal medicines, to elucidate analytical marker [2]. The major

highlights of plant derived drugs are their efficacy, low cost and selectively without or with little side effects.

Plumbago zeylanica Linn belonging to family Plumbaginaceae is native to the temperate regions of Europe and western and central Asia, east to the Himalaya and eastern Siberia, and America. It is an herbaceous perennial plant growing to 10–60 cm (rarely to 100 cm) tall. The leaves are in opposite pairs, simple, 2–6.5 cm long and 1–2 cm broad, hairy, and with a coarsely serrated margin. The flowers are pale purple (occasionally white or pink), in clusters on the stem, each flower 3–4 mm long. The plant is widely distributed throughout India and leaves of the plant are extensively used in traditional system of medicine for various ailments like carminative, digestive, expectorant, cardiogenic, diuretic, dentifrice, jaundice, hepatalgia, inflammation of liver, peptic ulcer, diarrhoea, bronchitis and skin diseases [11-14]. Antioxidants are molecules that can safely interact with free radicals and terminate their chain reaction before vital molecules are damaged [8]. Lack of antioxidants facilitates the development of degenerative disorders, cardiovascular diseases and cancer. One possible solution to this problem may be provided by the use of natural antioxidant compounds present in plant sources [9], which can be useful in protocols preventive medicine. Most synthetic antioxidants, such as Butylated

Hydroxytoluene (BHT) and Butylated Hydroxyl Anisole (BHA), which are widely used in the food industry, may have more damaging than beneficial effects as they could be responsible for liver damage and carcinogenesis [10].

For this reasons, interest in the use of natural antioxidants has increased now a days. The purpose of this study was to investigate and elucidate the phytochemical antioxidant and phenolic content in Leaves extracts of *P. zeylanica*.

2. MATERIALS AND METHODS

2.1. Reagents

Folin-Ciocalteu reagent and catechol were purchased from Merck Pvt. Ltd, India. 2,2-diphenylhydrazyl (DPPH), Thiobarbituric Acid (TBA), Trichloroacetic Acid (TCA), ascorbic acid and quercetin were obtained from Himedia lab, India. All other chemicals were of analytical grade.

2.2. Collection and preparation of plant material

P. zeylanica Leaves were cleaned, chopped, shade dried and powdered. Dried powder (50 g) was soxhlet extracted separately with 400 ml of acetone and methanol for 48 h. Both acetonetic and methanolic extracts were concentrated under reduced pressure using a rotary evaporator and then kept either refrigerated at 4 °C, or at -20 °C for long term storage.

2.3. Preliminary phytochemical screening

2.3.1 Detection of alkaloids

A drop of Mayer's reagent was added to a few milliliter of filtrate, by the side of the test tube. Formation of a white creamy precipitate indicated that the test was positive.

2.3.2. Detection for saponins

About 0.2 g of the extracts were shaken with 5 ml of distilled water and then heated to boil. The appearance of creamy mass of small bubbles (frothing) showed the presence of saponins.

2.3.3. Detection for tannins

Small quantities of extracts were mixed with water and heated in a water bath. The mixtures were filtered and ferric chloride was added to the filtrates. The formation of a dark green solution indicated the presence of tannins.

2.3.4. Detection for flavonoids

The extracts (0.5 ml) were dissolved in methanol. After the addition of a few fragments of Magnesium ribbon to the extracts followed by concentrated hydrochloric acid, the appearance of pink color indicated the presence of flavonoids.

2.3.5. Detection of Steroids

About 2 ml of acetic anhydride was added to 0.5 g of the extract of each with 2 ml of H₂SO₄. The color changed from violet to blue or green in some samples indicating the presence of steroids.

2.3.6. Detection of Glycosides

The extract was hydrolyzed with HCl solution and neutralized with NaOH solution. A few drops of Fehling's solution A and B for

minutes. An orange red precipitate indicates the presence of reducing sugars.

2.3.7. Detection of Phenols

To 1 ml of aqueous solution of plant extract, 2 ml of distilled water followed by few drops of 10% aqueous FeCl₃ solution are added. Formation of blue or green precipitate indicates the presence of phenols.

2.4. Evaluation of in vitro antioxidant activity

The in vitro antioxidant activity of the acetone and methanolic extracts of *P. zeylanica* Leaves was measured by the following assays:

2.4.1. Determination of total phenolic compounds in the extracts

The amount of total phenolic compounds in the extracts was determined using the Folin-Ciocalteu method [11, 12, 13]. Briefly, 0.5 ml of the sample were pipetted into a 10 ml volumetric flask containing 0.5 ml of Folin-Ciocalteu's reagent, 5 ml of distilled water and 1.5 ml of Na₂CO₃ solution (20% by weight), and the volume was made up to 10 ml with distilled water. During the oxidation of phenolic compounds, the phosphomolybdic and phosphotungstic acids contained in the Folin-Ciocalteu's reagent were reduced to blue-colored molybdenum and tungsten oxides. After 2 h, the absorbance of the blue coloration was measured at 765 nm against a blank sample. Measurements were compared to a standard curve prepared with gallic acid solutions, and expressed per

weight of dry extract. The total phenolic content was expressed as g gallic acid equivalent (GAE)/100 g dry weight. All samples were analyzed in triplicate.

2.4.2. Determination of total flavonoid content in the extracts

The total flavonoid content was determined colorimetrically by aluminum chloride method [12, 13]. Briefly, 0.5 ml solution of each plant extracts in methanol were separately mixed with 1.5 ml of methanol, 0.1 ml of 10% aluminum chloride, 0.1 ml of 1 M potassium acetate, and 2.8 ml of distilled water, and left at room temperature for 30 min. The absorbance of the reaction mixture was measured at 415 nm with a Hitachi UV/Visible spectrophotometer (CA, USA). Total flavonoid contents were calculated as quercetin equivalently from a calibration curve. The calibration curve was determined by preparing quercetin solutions at concentrations 12.5 to 100 mg/ml in methanol and expressed as milligrams of quercetin equivalents (QE/g of dry extract).

2.4.3. Evaluation of total antioxidant capacity

The total antioxidant capacity was measured by a spectrophotometric method [14]. At different concentration ranges, extract were dissolved in water and combined in eppendorf tubes with 1 ml of reagent solution (0.6 M H₂SO₄, 28 mM sodium phosphate, 4 mM ammonium molybdate). The tubes were incubated for 90 min at 95 °C. The mixture

was cooled to room temperature and the absorbance was read at 695 nm against a blank control. Ascorbic acid equivalents were calculated using standard graph prepared with various concentrations of ascorbic acid.

2.4.4. 2,2-diphenyl-1-picrylhydrazyl radical-scavenging activity

The antioxidant activity of the acetone and methanolic extracts of *P. zeylanica* Leaves was measured in terms of hydrogen donating or radical scavenging ability using the stable radical 2,2-diphenyl-1-picrylhydrazyl (DPPH). Reduction of the DPPH radical was determined by the decrease in its absorbance at 517 nm [15]. Ascorbic acid was used as standard reference.

2.4.5. Determination of reducing power

As the antioxidant activity of the extracts would be also manifested through their reducing power, the Fe³⁺ → Fe²⁺ transformation assay was established as a measure of reducing capacity. The reducing power of the *P. zeylanica* extracts was determined by the method of Oyaizu as previously described [16]. Extracts were first dissolved into 1 ml distilled water and then mixed with 2.5 ml 0.2 M phosphate buffer (pH 6.6) and 2.5 ml 1% potassium ferricyanide [K₃Fe (CN)₆]. The mixture was incubated at 50 °C for 20 min. Then, 2.5 ml of 10% TCA were added to the mixture, followed by centrifugation of 3000 rpm for 10 min. The upper layer (2.5 ml) was mixed with 2.5 ml distilled water and 0.5 ml 0.1 % FeCl₃, then

its absorbance at 700 nm with a Hitachi UV/Visible spectrophotometer. Ascorbic acid was used as a reference antioxidant compound.

2.4.6. Assay of hydroxyl radical-scavenging activity

The inhibitory effect of the extracts to prevent the degradation of deoxyribose by Fe³⁺ ions in the presence of H₂O₂-EDTA-ascorbate was determined in hydroxyl radical scavenging assays [17]. To 0.5ml of plant extract were mixed with reaction mixture containing FeCl₃ (100 µM), EDTA (104 µM), H₂O₂ (1 mM) and 2-deoxy-D-ribose (2.8 mM) in 1 ml final reaction volume made with potassium phosphate buffer (20 mM, pH 7.4) and were incubated for 1 h at 37 °C. The mixture was then heated at 95 °C in water bath for 15 min, followed by the addition of 1 ml each of TCA (2.8%) and TBA (0.5% TBA in 0.025 M NaOH containing 0.02% butylhydroxyanisole (BHA)). Finally the reaction mixture was cooled on ice and centrifuged at 5000 rpm for 15 min. The absorbance of the supernatant was measured at 532 nm. All readings were corrected for any interference from brown color of the extract or antioxidant by including appropriate controls. The negative control without any antioxidant was considered 100% deoxyribose oxidation. The percent hydroxyl radical scavenging activity of test sample was determined accordingly in comparison with the negative control. Ascorbic acid was taken as the

positive control

2.4.7. Statistical analysis

For display of data, points on a graph represent a mean ± standard deviation (SD; error bars) for an experimental group or observation.

3. RESULTS

3.1. Determination of plant yield

The percentage yield of *P. zeylanica* Leaves extract in methanol and acetone were analyzed. Among this, methanolic extract gave the highest yield (16.83%) while the yields were lower for the acetone (14.31%), ethanol (9.25%), petroleum ether (4.14%), and water (6.35%) extracts, respectively. Since both the methanolic and acetone extracts of *P. zeylanica* Leaves exhibited the highest yields, they were selected for the phytochemical screening.

3.2. Preliminary phytochemical analysis of *P. zeylanica*

The acetone and methanolic extracts of *P. zeylanica* leaves extract subjected for preliminary phytochemical screening revealed the presence of various phytochemical constituents as given in **Table 1**. It became evident from these results that methanolic extracts (EPZE) contained all analyzed phytochemicals and appeared richer when compared with the acetonetic Leaves extracts. These phytochemical compounds are known to support bioactive activities in medicinal plants and thus maybe responsible for the biological activities of the *P. zeylanica*

extracts used in this study.

3.3. Evaluation of in vitro antioxidant assay

3.3.1. DPPH radical scavenging activity

EPZE had substantial DPPH radical scavenging activity compared to ascorbic acid, and this activity increased in parallel to the increase in extract concentration, as shown in **Figure 1**. The IC₅₀ value of the EPZE extract was 79.1 µg/ml while that for reference ascorbic acid was 4.2 µg/ml. These results signify that the extract had distinct free radical scavenging activity in comparison to ascorbic acid.

3.3.2. Hydroxyl radical scavenging activity

EPZE extracts had also detectable hydroxyl radical scavenging activity compared to quercetin, and this activity increased proportionately to the increase in concentration of the extract, as shown in **Figure 2**. The estimated IC₅₀ values of EPZE and quercetin were 78.2 µg/ml and 20.9

µg/ml respectively. An increase in the reaction mixtures absorbance indicates an increase in the reducing capacity of the sample.

3.3.3. The reducing power potentials of the extract

Figure 3 shows the reducing power potential of EPZE in comparison with standard ascorbic acid at 700nm. The extract was found to have marked reducing capacity. The results illustrate that there was an increase in reducing power of the plant extract as the EPZE concentration increased in the assay.

3.3.4. Total phenolic contents, flavonoid contents and total antioxidants

Table 2 shows that the total phenolic content of *P. zeylanica* Leaves EPZE extract is 116.6±32.08 mg/g, whereas the total flavonoid content is 39.2±640 mg/g and the total antioxidant is 683±885 mg/g, which may specify with the previous studies [16, 18]

Table 1: Phytochemical screening of *P. zeylanica* Leaves extracts

Constituents	Acetone extract	Methanolic extract
Steroids	+	+
Alkaloids	+	+
Tannins	-	+
Glycosides	-	+
Phenols	-	+
Flavonoids	-	+
Saponins	+	+

“+” and “-” indicate the presence or absence of the constituents indicated




Figure 1: Percentage inhibitions of DPPH free radical activity of EPZE compared with standard antioxidant ascorbic acid. The data shown are the mean from three independent experiments. Each value is the mean±SD of three determinations




Figure 2: Percentage inhibition of hydroxyl radical scavenging activity of EPZE compared with quercetin as antioxidant standard. The data shown are the mean from three independent experiments. Each value is the mean±SD of three determinations




Figure 3: Reducing power of EPZE compared with ascorbic acid as antioxidant standard. The data shown are the mean from three independent experiments. Each value is the mean±SD of three determinations

Table 2: *In vitro* total phenolic and flavonoid contents and antioxidant activity of EPZE

Material	Total phenolic content (mg GAE/g dry extract)	Total flavonoid content (mg QE/g dry extract)	Total antioxidant (mg ascorbic acid/g dry extract)
EPZE	116.6±32.08	39.2±640	683.8±885

The data shown are the mean from three independent experiments.
Each value is the mean±SD of three determinations

4. DISCUSSION

Phytochemicals are defined as bioactive non-essential nutrients from plants, which possess a variety of human health effects such as possessing antioxidant and anti-carcinogenic activity [19]. Results on the percentage yield suggest that the methanolic extract of *P. zeylanica* Leaves (EPZE) was better than the acetonic extract. As a result EPZE was chosen for further studies.

The phytochemical analysis conducted on EPZE revealed the presence of alkaloids, flavonoids, saponins, glycosides and tannins. Alkaloids have been associated with medicinal uses for centuries and one of their common biological properties is their cytotoxicity [20], and their presence in this plant tend to increase the risk of poisoning by the plant. Flavonoids have been shown to exhibit their actions through effects on membrane permeability and by inhibition of membrane-bound enzymes, such as the ATPase and phospholipase A2 [21], a property that may explain the mechanism of antioxidative action of EPZE. The presence of flavonoids in EPZE may be responsible for its antioxidant activity. The antioxidant capacity of flavones is attributed to the high reactivity of the hydroxyl substituent, with

the number of hydroxyl groups on the B-ring being correlated with reactive oxygen species (ROS) scavenging capability [22, 23]. Also, the plant extract was revealed to contain saponins known to produce cytotoxic effect. The presence of these phenolic and non-phenolic compounds in these plant extracts contributed to their antioxidant properties and thus their usefulness of these plants in herbal medicament.

The *in vitro* antioxidant activity of EPZE indicates that the plant is potentially active. This suggests that the plant extracts contain compounds that are capable of donating hydrogen to a free radical in order to remove electrons responsible for the radicals reactivity. Though the DPPH radical scavenging activity of the EPZE extract was lower than that of the ascorbic acid reference, the study showed that EPZE have potent proton-donating ability and therefore could serve as free radical inhibitor. The reducing power of the EPZE extract was evaluated by the transformation of Fe³⁺ to Fe²⁺ through electron transfer ability, which serves as a significant indicator of its antioxidant activity. The reductive activity of the extract and that of the reference drug were augmented in parallel to their increasing concentrations, as

confirmed by escalating absorbance values at 700 nm. The antioxidant activity of the EPZE extract was found to be substantial when compared with the standard ascorbic acid standard used in this study.

Findings from this study showed that the antioxidant activity observed with the EPZE extract agrees well with the amount of phenolic constituents detected in the extract. The present data suggest that EPZE can effectively scavenge ROS, including hydroxyl radical as well as other free radicals, under *in vitro* conditions, and can be a potential source of natural antioxidant that could be of great importance for the treatment of radical-related diseases.

5. CONCLUSION

At present, a desirable portion of the global plant population still persist unexplored, and it seems reasonable to suggest that they hold the potential of the innovation for safer and more active new drugs. It is well known that medicinal plants are rich in antioxidants and can be used in the treatment of wide variety of diseases. Our findings show that the antioxidant activity of *P. zeylanica* extracts can be attributed to the pharmacological actions of alkaloids, saponins, flavonoids, tannins, glycosides, phenols and steroids present in the crude extract. These data strongly support the possible utility of these extracts in disease prevention and treatment.

REFERENCES

[1] Kumpulainen JT and Salonen JT: Natural

Antioxidants and Anticarcinogens in Nutrition. Health and Disease. The Royal Society of Chemistry, UK 1999; 178-187.

[2] Cook NC and Samman S: Flavonoids-chemistry, metabolism, cardioprotective effects, and dietary sources. *Nutritional Biochemistry* 1996; 7: 66-76.

[3] Deepika Gupta, Sonia Mann, Isha Jain and Rajinder K. Gupta: Phytochemical, nutritional and antioxidant activity evaluation of fruits of *Ziziphus nummularia burm F.* *International Journal of Pharma and Bio sciences* 2011; 2 (4): 629-638.

[4] Kerry NL and Abbey M: Red wine and fractionated phenolic compounds prepared from red wine inhibit low density lipoprotein oxidation *in vitro*. *Atherosclerosis* 1997; 135:93-102.

[5] Halliwell B and Gutteridge JMC: Formation of thiobarbituric acid reactive substances from deoxyribose in the presence of iron salts: the role of superoxide and hydroxyl radicals. *FEBS Lett* 1981; 128:347-352.

[6] Gulcin I: Antioxidant activity of food constituents, an overview. *Arch Toxicol* 2012, 86:345-391.

[7] Gocer H and Gulcin I: Caffeic acid phenethyl ester (CAPE): correlation of structure and antioxidant properties. *Int J Food Sci Nutr* 2011; 62:821-825.

[8] Nunes PX, Silva SF, Guedes RJ and Almeida S: Biological oxidations and

- antioxidant activity of natural products, Phytochemicals as nutraceuticals - Global Approaches to Their Role in Nutrition and Health 2012.
- [9] Djeridane A, Yousfi M, Nadjemi B, Boutassouna D, Stocker P and Vidal N: Antioxidant activity of some Algerian medicinal plants extracts containing phenolic compounds. *Food Chem* 2006; 97:654–660.
- [10] Wannas WA, Mhamdi B, Sriti J, Jemia MB, Ouchikh O, Hamdaoui G, Kchouk ME and Marzouk B: Antioxidant activities of the essential oil and methanol extracts from *myrtle* (*Myrtus communis* var. *Italica* L.) leaf, stem and flower. *Food Chem Toxicol* 2010; 48:1362–1370.
- [11] Sola AV: *Indian medicinal plants*. Orient Longman Private Ltd. 1995; 4: p. 15.
- [12] Kiritikar KR and Basu BD: *Indian medicinal Plants*, Allahabad, India: Lalit Mohan Basu 1998;2: p. 1982.
- [13] Chopra RN and Chopra IC: *Indigenous drugs of India*. Calcutta India, Academic Publishers 1994; 2: p. 196.
- [14] Khare CP: *Encyclopedia of Indian Medicinal plants*. Springer-Verlag Berlin Heidelberg, 2004: 309.
- [15] Verma SM, Arora H and Dubey R: Anti-Inflammatory and Sedative –Hypnotic Activity of The Methanolic Extract of The Leaves of *Plumbago zeylanica* . *Anc Sci Life* 2003; 23(2): 95–99.
- [16] Kowti R, Vishwanath S, Shivakumar SI, Vedamurthy J and Abdul NK: Hepatoprotective and Antioxidant Activity of Ethanol Extract of *Plumbago zeylanica* Leaves Against Carbon Tetrachloride Induced Hepatic Damage In Rats. *International Journal of PharmTech Research* 2013; 5(2): 426-430.
- [17] Coutinho HD, Costa JG, Lima EO, Falcao-Silva VS and Siqueira-Júnior JP: Potentiating effect of *Plumbago zeylanica* and chlorpromazine in the resistance to aminoglycosides of methicillin-resistant *Staphylococcus aureus*. *In vivo*. 2009 Mar-Apr;23(2):287-9.
- [18] Kanjanapothi D, Smitasiri Y, Pathong A, Taesotikul T and Rathanapanone V: Postcoital antifertility effect of *Plumbago zeylanica* . *Contraception* 1981; 24:559-567.
- [19] Marta C, Teixeira D and Glyn MF: Anticandida activity of Brazilian medicinal plants. *J Ethnopharmacol* 2005; 97: 305-311.
- [20] Ganesh CJ and Manjeshwar SB: Influence of the leaf extract of *Plumbago zeylanica* Linn. (mint) on the survival of Mice exposed to different doses of Gamma Radiation. *Strahlenther Onkol* 2002; 178: 91-8.
- [21] Akhtar HM, Anzar AK, Dar GH and Khan ZS: Ethnomedicinal uses of some plants in the Kashmir Himalaya. *Indian Journal of Traditional Knowledge* 2011; 10(2): 362-

366.

- [22] Towseef AB, Gaurav N and Masood M1: Study of Some Medicinal Plants of the Shopian District, Kashmir (India) With Emphasis on Their Traditional use by Gujjar and Bakerwal Tribes. Asian Journal of Pharmaceutical and Clinical Research 2012; 5(2).
- [23] Iyengar MA: Study of Crude Drugs. Manipal Power Press, Manipal, India, Edition 8th. 1995.