



**International Journal of Biology, Pharmacy  
and Allied Sciences (IJBPAS)**

*'A Bridge Between Laboratory and Reader'*

[www.jibpas.com](http://www.jibpas.com)

---

---

## EFFECT OF NORDIC WALKING AND TREADMILL WALKING ON CARDIORESPIRATORY FITNESS IN TYPE 2 DIABETES PATIENTS

SAIKIA D (PT)<sup>1</sup>, PATEL D (PT, PhD)<sup>2</sup> AND PATEL GJ (PT, PhD)<sup>3</sup>

- 1: MPT Student, Ahmedabad Physiotherapy College, Parul University, bopal-gnuma road,  
Ahmedabad, Gujarat
- 2: Associate Professor, Ahmedabad Physiotherapy College, Parul University. Ahmedabad, Gujarat
- 3: Principal, Ahmedabad physiotherapy College, Parul university, Parul University. Ahmedabad,  
Gujarat

\*Corresponding Author: Dr. Diyashree Saikia: E Mail: [yada.divya@gmail.com](mailto:yada.divya@gmail.com)

Received 24<sup>th</sup> Aug. 2023; Revised 25<sup>th</sup> Oct. 2023; Accepted 20<sup>th</sup> Jan. 2024; Available online 1<sup>st</sup> Nov. 2024

<https://doi.org/10.31032/IJBPAS/2024/13.11.8434>

### ABSTRACT

**Background & Purpose:** Diabetes mellitus is a prominent cause of mortality and disability worldwide. Diabetes mellitus is a condition of abnormal protein, lipid, and carbohydrate metabolism caused by inadequate insulin production and diminished sensitivity of the tissues to the metabolic effects of insulin. Increasing cardiorespiratory fitness play an important protective role in T2DM. Aerobic exercises has been shown to improve cardiorespiratory fitness in T2DM patients. The study focuses on the effectiveness of Nordic walking versus treadmill walking to improve cardiorespiratory fitness in type 2 diabetes patients.

**Methodology:** 40 type 2 diabetes patients with age group between 45-65 years were included. They were randomly allocated in 2 groups. Group A received Nordic walking program and Group B received treadmill walking program. In this study blood sugar was measured by fasting blood sugar, cardiorespiratory fitness was measured by 6-minute walk test and rockpot fitness walking test, quality of life was measured by SF36 health questionnaire before and after 8 weeks of intervention.

**Result:** Result were statistically analyzed using t test using SPSS 26. There was a significant improvement at  $p < 0.05$  in group A and group B. In between Group analysis there was more significant improvement in FBS, 6MWT, ROCKPOT and SF36 at  $p < 0.001$  in Group A than Group B.

**Conclusion:** This study showed that there is a positive effect of Nordic walking program and treadmill walking program to increase cardiorespiratory fitness in type 2 diabetes patients. But Group A showed greater significant difference as compared to Group B. Nordic walking program is more favorable when compared to treadmill walking program to improved cardiorespiratory fitness, blood sugar and health related quality of life.

**Keywords:** Type 2 diabetes, cardiorespiratory fitness, blood sugar, quality of life, Nordic walking, treadmill walking, FBS, 6MWT, ROCKPOT, SF36

## INTRODUCTION:

Diabetes is a group of metabolic diseases characterized by hyperglycemia due to defects in insulin secretion, insulin action, or both [1]. Diabetes mellitus is a syndrome of impaired carbohydrate metabolism, lipids and proteins caused by tissues decreased sensitivity to insulin metabolic effects [2]. The two broad type of diabetes mellitus are type 1 DM and type 2 DM. Both types of diabetes are caused by abnormal glucose homeostasis as the pathogenic processes progress. Type 1 DM is the result of complete or near-total insulin deficiency. Type 2 DM is characterized by high blood glucose, insulin resistance and relative insulin deficiency [3]

Diabetes affects both central and peripheral nervous systems. From every aspect of the nervous system and its supporting structures, from the cerebral cortex to the effector organs, may be involved with the complications of diabetes. These are caused by direct effect of hyperglycemia or hypoglycemia. Hyperglycemia induced metabolic derangements; neurochemical

alterations; serum lipid changes; and vascular, coagulation, and thrombotic abnormalities. Diabetic peripheral neuropathies are one of the most common complications of diabetes. Whether selective and pure nerve fiber damage occur in this form of polyneuropathy is still unclear [4, 5].

The overexpression of these cytokines contributes to increased inflammation and lipid accumulation, which have a effect on blood vessels and can lead to the development of endothelial dysfunction, MI and cardiomyopathy. Patients with DM also have decreased adiponectin production which may lead to increase in atherosclerotic plaque [6, 7].

Type 2 diabetes have physiologic exercise limitations and decreased cardiorespiratory fitness (approximately 20% lower maximal oxygen uptake i.e VO<sub>2</sub> max) when compared with those without diabetes that leads to decreased cardiorespiratory fitness and increased cardiovascular mortality [8]. Diabetes mellitus is a chronic disorder,

represents a heterogeneous group of disorders that have hyperglycemia as a common feature, and if left uncontrolled or mismanaged, the physical burden, emotional consequences, and activities of daily living are affected leading to a poor quality of life (QoL) of the individual [9].

Physical activity has been stated as one of the most important factors in the treatment of T2 DM and to improve and maintain the quality of life of the affected individual [2, 9]. Exercise training is associated with lowering blood pressure, improving insulin sensitivity, and glucose control and improving lipoprotein profile [8]. Exercise moves sugar into the muscles for storage and promotes an immediate increase in insulin sensitivity which last for 2-48 hours, depending on exercise [10]. One study found that 60 minutes of cycling on a machine at a moderate pace increased insulin sensitivity for 48 hours [11].

Cardiorespiratory fitness (CRF) also referred to as aerobic capacity is one of the most important components of physical fitness. Maximum oxygen consumption (VO<sub>2</sub>max) is considered to be the most widely accepted measure of CRF giving a baseline estimate of one's heart and lung capacity and can be used to follow the progress of daily physical exercise [12]. Higher levels of

cardiorespiratory fitness (CRF) have been shown to be associated with higher levels of health related quality of life [13].

Moderate to high levels of cardiorespiratory fitness also play a protective role in diabetes as individuals with low cardiorespiratory fitness are more likely to be insulin resistance [14]. CRF reflects one's capacity to achieve a certain exercise performance standard and increases skeletal muscle mass, improves autonomic function, decreases secretions of pro-inflammatory cytokines, and modifies the metabolic risk factors for type 2 diabetes [15].

Aerobic exercise training increases cardiorespiratory fitness in type2 diabetes patients [16]. Aerobic exercise is known to manage glycaemic control, cardiovascular risk factors and has also beneficial effects for metabolic profile in patients with T2DM. Aerobic exercise increases skeletal muscle capitalization and blood flow, muscular GLUT4 levels, hexokinase, and glycogen synthase activities [17]. In a meta-analysis study, aerobic exercise interventions of 8 weeks or more has been shown to be effective in improving cardiorespiratory fitness [18, 19].

Nordic walking is a form of physical activity requiring the use of specially designed poles, with which the force is pushed away

from the ground, resembling the Nordic Skiing style. The main advantage of the NW technique is involvement of muscles not used during standard walking, which results in higher energy expenditure. It was found that during NW, oxygen consumption is about 18–25% higher compared to walking without poles at the same speed. The NW technique makes this possible and the importance of maintaining high-intensity exercise and a low rate of perceived exertion [20].

In a recent studies shows that Nordic walking program is effective in improving BW control, body composition, muscular flexibility and VO<sub>2</sub> max levels in T2DM patients [19]. In one another article confirms that Nordic walking activities for 4 months have improved sleep quality and the overall increase of physical activities, which represents a significant element of treatment and preventive measures for patients of Type II Diabetes Mellitus [21].

#### **NEED OF STUDY**

Increasing cardiorespiratory fitness play an important protective role in T2DM. Aerobic exercises has been shown to improve cardiorespiratory fitness in T2DM patients according to different studies.

Among them, treadmill walking and Nordic walking both as individual have been proven for improving cardiorespiratory fitness. Both

these walking involves large muscle group of lower limbs as well as muscular work of upper limbs and is able to increase insulin sensitivity and reduce many cardiovascular risk factors in T2DM.

But there is no studies comparing the effects of this both techniques in T2DM patients. Hence this study will be carried out to compare the effect of Nordic walking and treadmill walking to determine which of this intervention is better in order to improve cardiorespiratory fitness among T2DM patients.

#### **AIM OF THE STUDY**

Aim of the study is to compare the effects of Nordic walking and treadmill walking on cardiorespiratory fitness in type 2 diabetes patients.

#### **MATERIAL AND METHODOLOGY**

Following ethical clearance number IECHR – SAINATH HOSPITAL/AHMC/50, an individual who met the inclusion criteria and fulfills initial evaluation were enrolled in this research. The consent form was filled once all of the participants had been informed about the study. They were distributed and assigned to either Group A or Group B, with 40 patients within each group.

**INCLUSION CRITERIA:** Both male and female patients between age group of 45-65 years, FBS:126-250 mg / dl, PPBS: 11.1 –

13.9 mmol/L, Non-smokers and non-alcoholic subjects, Willingness to participate in the research. **EXCLUSION CRITERIA:** Presence of any severe cardiovascular diseases, Presence of any musculoskeletal disorders, any medication that restrict them to participate, Uncontrolled hypertension, Central and peripheral nervous system disorders, diabetic retinopathy, myopathy, Inability to concentrate, follow direction, Rheumatological disease, Impaired knee flexion of <90-degree **OUTCOME MEASURE:** Fasting blood sugar, 6 minutes' walk test, Rock pot fitness walking test, SF 36 questionnaire

There are 2 groups, Group A = Nordic walking program

Group B = Treadmill walking program

Pre-assessment: Subject in both Groups were assessed by using fasting blood sugar test, 6-minute walk test, rockpot fitness walking test and SF36 questionnaire before starting the treatment.

#### **Procedure of intervention:**

All subjects were instructed to report immediately if they feel breathlessness, headache, dizziness or any other symptoms which are indicative for stopping exercise. Blood pressure, heart rate and respiratory rate were taken in supine at the beginning of exercise.

Throughout the duration of the exercises, blood pressure and respiratory, heart, and oxygen saturation rates were monitored with heart rate reserve maintained at moderate intensity of 40–65 % calculated by  $220 - \text{patient's age}$ .

#### **Group A (Nordic walking):**

Frequency – 3 times a week for 8 weeks  
Duration- 60 min

Warm up- (10-15) min, gentle mobility exercise. Low intensity aerobic walking, breathing exercise.

Exercise- 30 min in 1st 3 weeks with low intensity.

40 min in last 5 weeks with moderate intensity  
Cool down- (10-15) min, self-stretching of major muscle groups

#### **Group B (Treadmill walking):**

Frequency- 3 times a week for 8 weeks  
Duration- 60 min.

Warm up- (10-15) min, gentle mobility exercise. Low intensity aerobic walking, breathing exercise.

Exercise- speed 3.4mph with fixed inclination of 4.2 angle and Borg's score of 13 -14 for rate of perceived exertion.

#### **Statistical Analysis:**

The collected data were analysed using SPSS version 26.0. Paired and unpaired t-test are taken as the data followed normal distribution. According values they are allocated within

and between group comparison with p value <0.001.

## RESULT

In this study 40 patients were included. In group A there were 10 male and 10 female. In group B there were 11 male and 9 female.

The Mean score of Group A pre-FBS= 184.15, post-FBS= 145.70, Mean score of pre-6MWT= 408.20, post-6MWT= 627.25, Mean score of pre-Rockpot= 22.79, post-Rockpot= 40.55, Mean score of SF36: pre-physical functioning= 59.95, post physical functioning= 79.80, pre-physical health= 54.25, post-physical health= 75.30, pre-emotional problems= 0.00, post-emotional problems= 0.00, pre-energy/fatigue= 49.55, post-energy/fatigue= 79.95, pre-emotional well being= 50.10, post-emotional well being= 75.00, pre-social functioning= 46.90, post-social functioning=84.70, pre-pain= 51.20, post-pain= 75.50, pre general health= 46.75, post-general health=88.50.

pre and post mean data of FBS, 6MWT, ROCKPOT and SF36 of Group A. It suggest that there is significant difference in Group A (p<0.001)

The Mean score of Group B pre-FBS= 191.40, post-FBS= 164.05, Mean score of pre-6MWT= 413.60, post-6MWT= 578.10, Mean score of pre-Rockpot= 24.52, post-Rockpot=

40.98, Mean score of SF36: pre-physical functioning= 60.50, post physical functioning= 75.50, pre-physical health= 53.80, post-physical health= 72.65, Pre-emotional problems= 0.00, post-emotional problems= 0.00, pre-energy/fatigue= 50.05, post-energy/fatigue= 75.50, pre-emotional well being= 49.90, post-emotional well being= 70.85, pre-social functioning= 50.60, post-social functioning=79,90, pre-pain= 47.30, post-pain= 105.70, pre general health= 44.20, post-general health=76.95.

Pre-post data of FBS, 6MWT, ROCKPOT and SF36 in Group A and Group B were assessed. Paired t-test was used and p value is <0.001. As a result, there was a considerable difference between the two groups. But Group A showed more significance improvement than Group B.

For group A mean score of FBS = 145.70 and group B mean score of FBS = 164.05, for group A mean score of 6 MWT = 627.25 and group B mean score of 6 MWT = 578.10, for group A mean score of ROCKPOT = 40.55 and group B mean score of ROCKPOT = 40.98, for group A mean score of SF36, post physical functioning= 79.80, post-physical health= 75.30, post-emotional problems= 0.00, post-energy/fatigue= 79.95, post-emotional well being= 75.00, post-social

functioning=84.70, post-pain= 75.50, post-general health=88.50 and for group B mean score of SF36, post physical functioning=75.50, post-physical health= 72.65, post-emotional problems= 0.00, post-energy/fatigue=75.50, post-emotional well being=70.85, post-social functioning=79,90, post-pain= 105.70, post-general health=76.95. So, this shows Group A more effective rather than Group B. Group A and Group B intergroup significance  $p < 0.01$ . So, these

results reveal that there is a considerable difference between Group A and Group B. As a result, the null hypothesis is inapplicable to this study.

Sample t-test was used to analysis of intergroup data. After comparison between both the Groups, significantly greater improvement was observed in GROUP A. Statistically NORDIC WALKING PROGRAM is more effective to increase cardiorespiratory fitness in type2 diabetes patients.

**Table 1.1: Comparison of Pre and Post Mean In TreadmillWalking Program (Group B)**

Outcomes Measures	TREADMILL WALKING		P-VALUE
	Mean	SD	
PRE FBS (mg/dl)	191.40	13.76	P<0.001
POST FBS (mg/dl)	164.05	12.17	
PRE 6MWT (meter)	413.60	40.57	P<0.001
POST 6MWT (meter)	578.10	27.13	
PRE ROCKPOT (ml/kg/min)	24.52	8.80	P<0.001
POST ROCKPOT (ml/kg/min)	40.98	7.14	
pre physical functioning	60.50	5.70	P<0.001
post physical functioning %	75.50	3.78	
pre physical health	53.80	6.22	P<0.001
post physical health %	72.65	2.21	
Pre emotional problems	0.00	0.00	1.000
post emotional problems %	0.00	0.00	
Pre energy/fatigue	50.05	3.63	P<0.001
post energy/fatigue %	75.50	3.78	
Pre emotional well being	49.90	3.55	P<0.001
post emotional well being %	70.85	2.94	
pre social functioning	50.60	4.25	P<0.001
post social functioning %	79.90	3.92	
pre pain	47.30	2.64	P<0.001
post pain%	105.70	153.09	
pre general health	44.20	3.81	P<0.001
post general health %	76.95	5.86	

Table 1.2: Between Group Comparison

Outcomes Measures	TREADMILL WALKING		P-VALUE	NORDIC WALKING			P-VALUE (between group)
	Mean	SD		Mean	SD	P-VALUE	
pre physical functioning	60.50	5.70	P<0.001	59.95	6.18	P<0.001	0.870
post physical functioning %	75.50	3.78		79.80	3.65		0.003
pre physical health	53.80	6.22	P<0.001	54.25	6.22	P<0.001	0.764
post physical health %	72.65	2.21		75.30	3.57		0.018
pre emotional problems	0.00	0.00	1.000	0.00	0.00	1.000	1.000
post emotional problems %	0.00	0.00		0.00	0.00		1.000
pre energy/fatigue	50.05	3.63	P<0.001	49.55	5.61	P<0.001	0.989
post energy/fatigue %	75.50	3.78		79.95	3.50		0.002
pre emotional well being	49.90	3.55	P<0.001	50.10	5.51	P<0.001	0.763
post emotional well being %	70.85	2.94		75.00	3.46		0.001
pre social functioning	50.60	4.25	P<0.001	46.90	4.13	P<0.001	0.022
post social functioning %	79.90	3.92		84.70	3.74		P<0.001
pre pain	47.30	2.64	P<0.001	51.20	4.71	P<0.001	0.008
post pain %	105.70	153.09		75.50	3.36		0.004
pre general health	44.20	3.81	P<0.001	46.75	6.00	P<0.001	0.188
post general health %	76.95	5.86		88.50	2.48		P<0.001
PRE FBS (mg/dl)	191.40	13.76	P<0.001	184.15	13.60	P<0.001	0.62
POST FBS (mg/dl)	164.05	12.17		145.70	10.44		0.27
PRE 6MWT (meter)	413.60	40.57	P<0.001	408.20	32.44	P<0.001	0.05
POST 6MWT (meter)	578.10	27.13		627.25	34.77		0.29
PRE ROCKPOT (ml/kg/min)	24.52	8.80	P<0.001	22.79	8.37	P<0.001	0.76
POST ROCKPOT (ml/kg/min)	40.98	7.14		40.55	7.99		0.55

PHOTOGRAPHS:



Figure: 1.1



Figure: 1.2

---

## DISCUSSION

The impact of Nordic walking versus treadmill walking on cardiorespiratory fitness in type 2 diabetes patients is examined in this study.

In this study group A was given 60 minutes Nordic walking program. After intervention blood sugar, walking distance, Vo2 max and quality of life evaluated by FBS, 6MWT, ROCKPOT and SF36. The result of pre intervention mean score of FBS=184.15, 6MWT= 408.20, ROCKPOT = 22.79 and SF36= pre-physical functioning= 59.95, pre- physical health= 54.25, pre-emotional problems= 0.00, pre-energy/fatigue= 49.55, pre- emotional well being= 50.10, pre-social functioning= 46.90, pre-pain= 51.20, pre general health= 46.75. Post intervention mean score of FBS=145.70, 6MWT= 627.25, ROCKPOT= 40.55 and SF36=post physical functioning= 79.80, post-physical health= 75.30, post-emotional problems= 0.00, post-energy/fatigue= 79.95, post-emotional well being= 75.00, post-social functioning=84.70, post-pain= 75.50, post-general health=88.50.

According to result there is significant difference of FBS, 6MWT, ROCKPOT and SF36 within the group. So there is improvement in blood sugar, cardiorespiratory fitness and quality of life within the group.

Roberto pippi, *et al.* (2020) had done a study to assess the impact of Nordic walking program on obese adults with and without type 2 diabetes. 108 obese adults, 45-65 years aged female and male with or without diabetes mellitus were taken. Protocol was given for 3 months, with 90 min duration for 2 days per week. The intervention consists of 2 forms of supervised exercise programs (Nordic walking and gym based exercise program). They used BMI, blood pressure, FBS, Hba1c, blood chemistry, fitness variables, such as aerobic fitness (maximal oxygen uptake, VO2 max) and muscular flexibility were evaluated using the Rockport fitness walking test and the Bending test (executed from vertical and horizontal position), respectively as outcome measures. This study showed significant improvement in all the outcome measures and it provided evidence to support the Nordic walking program.

T. fritz, *et al* (2011) had done a study to assess the impact of Nordic walking program in overweight individuals with type 2 diabetes mellitus, impaired or normal glucose tolerance. 212 overweight individuals 57-64 years aged female and male with type 2 diabetes mellitus were taken. Protocol was given for 4 months with 5 hour duration per week. They used self reported physical activity and health related quality of life as outcome measures. This

study showed significant improvement in quality of sleep following 4 months of Nordic walking. Nordic walking can be introduced in a primary healthcare setting as a low cost mode of exercise that promotes weight loss and improved health satisfaction.

In group B was given 60 minutes treadmill walking program. After intervention blood sugar, walking distance, Vo2 max and quality of life evaluated by FBS, 6MWT, ROCKPOT and SF36.

The result of pre intervention mean score of FBS=191.40, 6MWT= 413.60, Rockpot=24.52 and SF36: pre-physical functioning= 60.50, pre-physical health= 53.80, pre-emotional problems= 0.00, pre-energy/fatigue= 50.05, pre-emotional well being= 49.90, pre-social functioning= 50.60, pre-pain= 47.30, pre general health= 44.20. Post intervention mean score of FBS=164.05, 6MWT= 578.10, Rockpot= 40.98 and SF36: post physical functioning= 75.50, post-physical health= 72.65, post-emotional problems= 0.00, post-energy/fatigue= 75.50, post-emotional well being= 70.85, post-social functioning=79.90, post-pain= 105.70, post-general health=76.95. According to result there is significant difference of FBS, 6MWT, ROCKPOT and SF36 within the group. So there is improvement in blood

sugar, cardiorespiratory fitness and quality of life within the group.

Paras arvindbhai parekh (2019) had done a study to assess the impact of treadmill exercise on blood glucose control in type 2 diabetes mellitus patients. 40 type 2 diabetes mellitus male patients, 45-55 years aged were taken. Protocol was given for 8 weeks with 40 minute duration, which include 5 minute of warm up, 30 minute treadmill walking followed by 5 minute of rest. They used FBS and PPBG level as outcome measures. This study showed significant decrease in FBS and PPBG level. Treadmill exercise is very useful for blood glucose control in addition to diet control and medicines. This article provide evidence to support treadmill walking program.

Sonill S. Maharaj *et al* (2014) had done a study to assess the impact of treadmill walking and rebound exercises on quality of life for patients with non-insulin dependent type 2 diabetes. 150 type 2 diabetes mellitus patients were taken. Protocol was given for 12 weeks with 30 min duration 3 times per week. They used SF36 to measure quality of life as outcome measure. This study showed significant improvement in quality of life after treadmill walking. Treadmill walking can be used to improve QoL for T2D patients and possibly reduce the side effects and co-morbidities associated with diabetic medication and diabetes.

Exercise is crucial for the treatment and prevention of type 2 diabetes since it aids in the management of the condition's associated lipid, glucose, and blood glucose control problems as well as the maintenance and reduction of weight. Aerobic exercise has been shown to lower the risk of glucose intolerance condition in people with non-insulin dependent diabetes. Walking is regarded as the finest workout since it works both the muscles in the upper and lower extremities.

Result of this study showed that Nordic walking program and treadmill walking program both the group individually statistically significant to improve blood sugar level, cardiorespiratory fitness and quality of life. But Nordic walking program showed more improved effect when compared to treadmill walking program. Nordic walking program showed more improvement because Nordic walking has a greater cardiometabolic efficacy likely due to the continuous active use of muscles of trunk, upper and lower limbs compared to treadmill walking.

This study proves that Nordic walking program and treadmill walking program improved cardiorespiratory fitness in type 2 diabetes patients. So from this result we confirmed that 8 weeks of Nordic walking program shows more improvement in cardiorespiratory fitness in type 2 diabetes

population while comparing both groups, group A showed more improved scores than group B.

### CONCLUSION

This study showed that there is a positive effect of Nordic walking program and treadmill walking program to increase cardiorespiratory fitness in type 2 diabetes patients. But Group A showed greater significant difference as compared to Group B. Nordic walking program is more favorable when compared to treadmill walking program to improved cardiorespiratory fitness, blood sugar and health related quality of life.

### REFERENCES:

- [1] Benkar AP, Kanase SB. Effect of aerobic exercises and resistance training on glucose levels in type 2 diabetic subjects. *Asian J Pharm Clin Res.* 2017 Jul 1;10(7):360.
- [2] Motahari-Tabari N, Shirvani MA, Shirzad-e-Ahoodashty M, Yousefi-Abdolmaleki E, Teimourzadeh M. The effect of 8 weeks aerobic exercise on insulin resistance in type 2 diabetes: a randomized clinical trial. *Global journal of health science.* 2015;7(1):115.
- [3] Parekh PA. Effect of Treadmill Exercise on Blood Glucose Control in Type-2 Diabetes Mellitus Patients. *International Journal of*

- Physiology. 2019 Jul 20;7(2):96-8.
- [4] Harati Y. Diabetes and the nervous system. Endocrinology and metabolism clinics of North America. 1996 Jun 1;25(2):325-59.
- [5] Kim RP, Edelman SV, Kim DD. Musculoskeletal complications of diabetes mellitus. Clinical diabetes. 2001 Jul 1;19(3):132-5.
- [6] Leon BM, Maddox TM. Diabetes and cardiovascular disease: Epidemiology, biological mechanisms, treatment recommendations and future research. World journal of diabetes. 2015 Oct 10;6(13):1246.
- [7] Martín-Timón I, Sevillano-Collantes C, Segura-Galindo A, del Cañizo-Gómez FJ. Type 2 diabetes and cardiovascular disease: have all risk factors the same strength?. World journal of diabetes. 2014 Aug 15;5(4):444.
- [8] Wahl MP, Scalzo RL, Regensteiner JG, Reusch JE. Mechanisms of aerobic exercise impairment in diabetes: a narrative review. Frontiers in endocrinology. 2018 Apr 18;9:181.
- [9] Maharaj SS, Nuhu JM. The effect of rebound exercise and treadmill walking on the quality of life for patients with non-insulin-dependent type 2 diabetes. International Journal of Diabetes in Developing Countries. 2015 Sep;35(2):223-9.
- [10] Borghouts LB, Keizer HA. Exercise and insulin sensitivity: a review. International journal of sports medicine. 2000 Jan;21(01):1-2
- [11] Magkos F, Tsekouras Y, Kavouras SA, Mittendorfer B, Sidossis LS. Improved insulin sensitivity after a single bout of exercise is curvilinearly related to exercise energy expenditure. Clinical Science. 2008 Jan 1;114(1):59-64.
- [12] Hingorjo MR, Zehra S, Hasan Z, Qureshi MA. Cardiorespiratory fitness and its association with adiposity indices in young adults. Pakistan Journal of Medical Sciences. 2017 May;33(3):659.
- [13] Sloan RA, Sawada SS, Martin CK, Church T, Blair SN. Associations between cardiorespiratory fitness and health-related quality of life. Health and Quality of Life Outcomes. 2009 Dec;7(1):1-5.
- [14] Loprinzi PD, Pariser G. Cardiorespiratory fitness levels and its correlates among adults with diabetes. Cardiopulmonary physical therapy journal. 2013 Jun;24(2):27.

- 
- [15] Qiu S, Cai X, Yang B, Du Z, Cai M, Sun Z, Zügel M, Michael Steinacker J, Schumann U. Association between cardiorespiratory fitness and risk of type 2 diabetes: A meta-analysis. *Obesity*. 2019 Feb;27(2):315-24.
- [16] Sigal RJ, Armstrong MJ, Bacon SL, Boule NG, Dasgupta K, Kenny GP, Riddell MC. Physical activity and diabetes. *Canadian journal of diabetes*. 2018 Apr 1;42:S54-63.
- [17] Yavari A, Najafipour F, Aliasgarzadeh A, Niafar M, Mobasser M. Effect of aerobic exercise, resistance training or combined training on glycaemic control and cardiovascular risk factors in patients with type 2 diabetes. *Biology of Sport*. 2012 Jan 1;29(2):135.
- [18] Boulé NG, Kenny GP, Haddad E, Wells GA, Sigal RJ. Meta-analysis of the effect of structured exercise training on cardiorespiratory fitness in Type 2 diabetes mellitus. *Diabetologia*. 2003 Aug;46(8):1071-81.
- [19] Pippi R, Di Blasio A, Aiello C, Fanelli C, Bullo V, Gobbo S, Cugusi L, Bergamin M. Effects of a supervised nordic walking program on obese adults with and without type 2 diabetes: The CURIA Mo. Centre experience. *Journal of functional morphology and kinesiology*. 2020 Sep;5(3):62.
- [20] Kantorowicz M, Szymura J, Szygula Z, Kusmierczyk J, Maciejczyk M, Wiecek M. Nordic Walking at Maximal Fat Oxidation Intensity Decreases Circulating Asprosin and Visceral Obesity in Women With Metabolic Disorders. *Frontiers in Physiology*. 2021 Sep 3:1457.
- [21] Upeniece I, Smite D, Arnis V. Nordic Walking for Patients of Type II Diabetes Mellitus: A Critical Review. *Education in a Changing Society*. 2014 May 29;1:108-12.
-